Anti-Human PLASMINOGEN ACTIVATOR INHIBITOR TYPE 1 (PAI-1)

(Polyclonal Rabbit)

Cat. No.: 1020A

Isotype: Mostly IgG.

Antigen: Purified PAI-1 from the HT 1080 fibrosarcoma cell line.

Titre: 1:100 to 1:10,000 depending on the application.

Cross-reactions: Reacts with Human PAI-1s from endothelial cells. No reactivity

with human PAI-2. Identity established by double diffusion with human PAI-1 isolated from human endothelial cells (HUVEC) culture and recombinant PAI-1 expressed in E.coli. No other proteins so far tested showed greater than 1% cross reaction.

Characteristics: Neutralises inhibitor's biological activity, recognises t-PA-PAI-1

and u-PA-PAI-1 complexes.

Presentation: 1ml unpurified serum and no preservatives.

Applications: Activity inhibition characterisation (especially coupled with reverse

fibrin autography) and RIA. Immunoprecipitation,

immunocytochemistry using frozen or paraffin-fixed sections, Western Immunoblotting, ELISA, fluorescence activated cell

sorting.

Storage: +4 °C for 1 month provided preservative is added For extended

storage, freeze in working aliquots at -20 °C. Avoid repeated

freezing and thawing.

FOR RESEARCH USE ONLY